
A trait-based approach to understand and predict the performance of arable annual mixed crops

Rémi Mahmoud¹, Noémie Gaudio^{*1}, Pierre Casadebaig¹, Xavier Gendre², Laurent Bedoussac^{1,3}, Guénaëlle Corre-Hellou⁴, Florian Fort⁵, Etienne-Pascal Journet^{1,6}, Isabelle Litrico⁷, Christophe Naudin⁴, and Cyrille Violle⁵

¹AGroécologie, Innovations, teRritoires (AGIR) – Institut National de la Recherche Agronomique : UMR1248 – Centre de Recherches INRA de Toulouse Chemin de Borderouge 31326 CASTANET TOLOSAN, France

²Institut de Mathématiques de Toulouse UMR5219 (IMT) – Centre National de la Recherche Scientifique, Université Paul Sabatier - Toulouse 3 – UPS IMT, F-31062 Toulouse Cedex 9, INSA Toulouse, F-31077 Toulouse, France UT1, F-31042 Toulouse, France UT2, F-31058 Toulouse, Téléphone : 05.61.55.67.90, France

³École Nationale Supérieure de Formation de l'Enseignement Agricole de Toulouse-Auzeville (ENSFEA) – ENSFEA – BP 226872 Route de Narbonne, 31326 Castanet Tolosan Cedex, France

⁴ESA - Ecole supérieure d'Agriculture d'Angers (ESA) – LEVA, INRA – Enseignement supérieur et recherche en agriculture, alimentation, territoires et marchés 55 Rue Rabelais, 49007 Angers – 02 41 23 55 55, France

⁵Centre d'Ecologie Fonctionnelle et Evolutive (CEFE) – Université de Montpellier : UMR5175, Centre National de la Recherche Scientifique : UMR5175 – 1919 route de Mende - 34293 Montpellier cedex 5, France

⁶Laboratoire des interactions plantes micro-organismes (LIPMO) – Centre National de la Recherche Scientifique : UMR2594 – Chemin de Borde-Rouge - BP 27 31326 CASTANET TOLOSAN CEDEX, France

⁷Unité de Recherche Pluridisciplinaire Prairies et Plantes Fourragères (P3F) – Institut National de la Recherche Agronomique – LUSIGNAN, France

Abstract

One of the challenges in low-input annual arable mixed crops (i.e. species and variety mixtures) is to determine which species / varieties go well together to improve overall system performance.

Theories coming from community ecology suggest that two ecological processes are particularly involved in mixed crops performance by improving resource use efficiency [1]. Niche complementarity can be quantified by the distance between target functional traits between two plants. Phenotypic plasticity can be quantified by the variance of target functional traits between cropping environments (including crop management). We aim to apply this ecological theory in an agronomical context by calculating these complementarity and plasticity metrics to predict crop yield of various mixed crops in different environments.

*Speaker

A plant trait database on wheat and pea mixtures was built, gathering ten experiments with various management practices over six years, localized in two French sites (Angers, Toulouse). Plant traits were measured dynamically (crop height, aerial biomass, nitrogen content) to quantify crop growth and development over key phenological stages and at harvest (yield, grain quality) to quantify crop performance.

Together with environmental (climate and soil) indicators we considered both trait by trait and multi-traits methods to analyze performance variability.

Litrico and Violle (2015). *Trends Plant Sci.* 20, 604-613.